

## Supplementary Information

### Design of Novel *N*-Myristoyltransferase Inhibitors of *Leishmania donovani* Using Four-Dimensional Quantitative Structure-Activity Relationship Analysis

*Letícia Santos-Garcia, Daniela R. Silva, Letícia C. Assis, Tamiris M. de Assis, Giovanna C. Gajo, Ítalo Antônio Fernandes, Teodorico C. Ramalho and Elaine F. F. da Cunha\**

*Departamento de Química, Universidade Federal de Lavras,  
PO Box 3037, 37200-000 Lavras-MG, Brazil*

**Table S1.** Calculated parameters of the Lipinski's Rule for the proposed molecules

Compound	logP	MW	n <sub>ON</sub>	n <sub>OHNH</sub>	n <sub>rotb</sub>	n <sub>violations</sub>
<b>A</b>	2.38	420.47	8	1	6	0
<b>B</b>	4.01	528.40	8	1	7	1
<b>C</b>	3.26	479.53	9	1	8	0

logP: partition-coefficient; MW: molecular weight; n<sub>ON</sub>: number of hydrogen bond acceptors; n<sub>OHNH</sub>: number of hydrogen bond donors; n<sub>rotb</sub>: number of rotatable bonds; n<sub>violations</sub>: number of violations.

\*e-mail: elaine\_cunha@dqi.ufla.br

