

Supplementary Information

Identification of Ethyl and *t*-Butyl Glyceryl Ethers Using Gas Chromatography Coupled with Mass Spectrometry

*Paula M. Veiga, Cláudia O. Veloso, Ayres G. Dias and Cristiane A. Henriques**

Instituto de Química, Universidade do Estado do Rio de Janeiro, 20550-900 Rio de Janeiro-RJ, Brazil




Figure S1. Mass spectrum of 3-ethoxy-1,2-propanediol (**1a**).




Figure S2. Mass spectrum of 2-ethoxy-1,3-propanediol (**1b**).




Figure S3. Mass spectrum of 1,3-diethoxy-2-propanol (**2a**).




Figure S4. Mass spectrum of 2,3-diethoxy-1-propanol (**2b**).




Figure S5. Mass spectrum of 1,2,3-triethoxypropane (**3**).




Figure S6. Mass spectrum of 3-*tert*-butoxy-1,2-propanediol (**4a**).




Figure S7. Mass spectrum of 2-*tert*-butoxy-1,3-propanediol (**4b**).




Figure S8. Mass spectrum of 1,3-di-*tert*-butoxy-3-propanol (**5a**).




Figure S9. Mass spectrum of 2,3-di-*tert*-butoxy-1-propanol (**5b**).