

## Using a Multivariate Approach to Compare Lipid Extraction Protocols from Microalgae *Scenedesmus* sp.

*Daiane F. Dall'Oglio,<sup>a,b</sup> Láiisse C. de Sousa,<sup>a</sup> Samuel A. A. de Sousa,<sup>a</sup> Marco A. S. Garcia,<sup>a</sup> Edymilaís S. Sousa,<sup>a</sup> Sidney G. de Lima,<sup>a</sup> Pelrry S. Costa,<sup>a</sup> Abhishek Guldhe,<sup>c</sup> Faizal Bux,<sup>c</sup> Edmilson M. de Moura<sup>a</sup> and Carla V. R. de Moura<sup>\*,a</sup>*

<sup>a</sup>*Departamento de Química, Universidade Federal do Piauí, 64049-550 Teresina-PI, Brazil*

<sup>b</sup>*Centro de Ciências Agrárias e Ambientais, Universidade Federal do Maranhão, 65500-000 Chapadinha-MA, Brazil*

<sup>c</sup>*Institute for Water and Wastewater Technology, Durban University of Technology, P.O. Box 1334, 4000 Durban, South Africa*

**Table S1.** ANOVA table evaluating the extraction time effect (2, 4 and 6 h) on the lipid percentage of the microalgae *Scenedesmus* sp.

Source	ANOVA				
	Sum of squares	Degrees of freedom	Mean square	<i>F</i>	<i>p</i>
Between group	2.18447	2	1.09223	3.04	0.1223
Within group	2.15353	6	0.35892		
Total	4.338	8			

\*e-mail: carla@ufpi.edu.br