

A New Eremophilane-type Sesquiterpene from the Phytopathogen Fungus
Lasiodiplodia theobromae (Sphaeropsidaceae)

Fátima M. Nunes,^a Maria da Conceição F. de Oliveira,^{*a} Ângela M. C. Arriaga,^a Telma L. G. Lemos,^a
Manoel Andrade-Neto,^a Marcos C. de Mattos,^a Jair Mafezoli,^b Francisco M. P. Viana,^c
Viviane M. Ferreira,^c Edson Rodrigues-Filho^d and Antônio G. Ferreira^d

^a Departamento de Química Orgânica e Inorgânica, Universidade Federal do Ceará, Campus do Pici,
CP 6044, 60455-970 Fortaleza-CE, Brazil

^b Curso de Farmácia, Universidade de Fortaleza, Av. Washington Soares, 1321,
CP 1258, 60811-341 Fortaleza-CE, Brazil

^c Laboratório de Fitopatologia, Embrapa – Agroindústria Tropical, Fortaleza-CE, Brazil

^d Departamento de Química, Universidade Federal de São Carlos, Rodovia Washington Luiz, km 235,
CP 676, 13565-905 São Carlos-SP, Brazil




Figure S1. ¹H NMR spectrum of **1** (400 MHz, CDCl₃).

*e-mail: mcfo@ufc.br




Figure S2. ^{13}C NMR spectrum of **1** (50 MHz, CDCl_3).




Figure S3. ^1H - ^1H COSY NMR correlation spectroscopy 2D NMR spectrum of **1** (400 MHz, CDCl_3).




Figure S4. ^1H - ^{13}C HSQC 2D NMR correlation spectroscopy of **1** (400 MHz/100 MHz, CDCl_3).




Figure S5. ^1H - ^{13}C HMBC 2D NMR correlation spectroscopy of **1** (400 MHz/100 MHz, CDCl_3).




Figure S6. APCIMS $[M+H]^+$ daughter ions spectrum of 1: (A) 20 eV (B) 10 eV.




Figure S7. nOe difference spectra for H-3, H-6 and H-4 in **1** (400 MHz, CDCl_3).