

Antiparasitic Activities of Acridone Alkaloids from *Swinglea glutinosa* (Bl.) Merr.

Djalma A. P. dos Santos,^a Paulo C. Vieira,^a M. Fátima das G. F. da Silva,^a
João B. Fernandes,^a Lauren Rattray^b and Simon L. Croft^b*

^aDepartamento de Química, Universidade Federal de São Carlos, CP 676, 13565-905 São Carlos-SP, Brazil

^bDepartment of Infectious and Tropical Diseases, London School of Hygiene & Tropical Medicine,
London, WC1E 7HT, UK




Figure S1. ¹H NMR spectrum of compound 1 (CDCl₃, 400 MHz).




Figure S2. ^{13}C NMR spectrum of compound 1 (CDCl_3 , 100 MHz).




Figure S3. HMBC spectrum of compound d 1 (CDCl_3).




Figure S4. HMQC spectrum of compound 1 (CDCl_3).








Figure S5. High resolution mass spectrum of compound 1.

Figure S6. ¹H NMR of compound 2 (CDCl₃, 400 MHz).Figure S7. ¹³C NMR of compound 2 (CDCl₃, 100 MHz).

Figure S8. HMBC spectrum of compound 2 (CDCl_3).Figure S9. HMQC spectrum of compound 2 (CDCl_3).





Figure S10. High resolution mass spectrum of compound 2.

Figure S11. ¹H NMR of compound 3 (CDCl₃, 400 MHz).




Figure S12. ^{13}C NMR of compound **3** (CDCl_3 , 100 MHz).




Figure S13. HMBC spectrum of compound **3** (CDCl_3).



Figure S14. HMQC spectrum of compound 3 (CDCl_3).

Figure S15. High resolution mass spectrum of compound 3.