

Supplementary Information

Organonickel(II) Complexes with Anionic Tridentate 1,3-Bis(azolylmethyl)phenyl Ligands. Synthesis, Structural Characterization and Catalytic Behavior

John Hurtado,^a Andrés Ibañez,^b René Rojas,^a Mauricio Valderrama^{*,a} and Roland Fröhlich^c

^aDepartamento de Química Inorgánica, Facultad de Química,
Pontificia Universidad Católica de Chile, Casilla 306, 6094411 Santiago, Chile

^bCentro para la Investigación Interdisciplinaria Avanzada en Ciencia de los Materiales (CIMAT),
Santiago, Chile

^cOrganisch Chemisches Institut der Universität Münster, Corrensstrasse 40,
48149 Münster, Germany




Figure S1. ¹H NMR (CDCl₃) spectrum of 2-bromo-1,3-bis(indazol-2-ylmethyl)benzene.




Figure S2. HMBC spectrum of 2-bromo-1,3-bis(indazol-2-ylmethyl)benzene.




Figure S3. HSQC spectrum of 2-bromo-1,3-bis(indazol-2-ylmethyl)benzene.




Figure S4. ^1H NMR (acetone) spectrum of $\text{NiBr}\{1,3\text{-bis(indazol-2-ylmethyl)phenyl-}N,C,N\}$.




Figure S5. HMBC spectrum of $\text{NiBr}\{1,3\text{-bis(indazol-2-ylmethyl)phenyl-}N,C,N\}$.