

Supplementary Information

Coprecipitation of Safrole Oxide with Poly(3-hydroxybutyrate-co-3-hydroxyvalerate) in Supercritical Carbon Dioxide

Ernandes T. Tenório-Neto,^a Expedito L. Silva,^a Thelma S. Pacheco Cellet,^a Elisangela P. Silva,^a Elton Francheschi,^b Lúcio C. Filho,^c Adley F. Rubira^a and Marcos H. Kunita^{*a}

^aDepartamento de Química, Universidade Estadual de Maringá,
Av. Colombo, 5790, 87020-900 Maringá-PR, Brazil

^bInstituto de Tecnologia e Pesquisa (ITP), Universidade Tiradentes,
Campus Farolândia, Av. Murilo Dantas, 300, 49032-490 Aracajú-SE, Brazil

^cDepartamento de Engenharia Química, Universidade Estadual de Maringá,
Av. Colombo, 5790, 87020-900 Maringá-PR, Brazil




Figure S1. ¹H NMR spectrum of safrole.




Figure S2. ¹H NMR spectrum of safrole oxide.





Figure S3. Calibration curve of safrole oxide in different solvents, using as reference λ_{\max} 287 nm. Insets show safrole oxide absorption, having as solvent (a) dichloromethane and (b) hydroalcoholic solution (62.5% ethanol).

*e-mail: mkunita@gmail.com