

Supplementary Information

Chemometric Study of the Antiproliferative Activity of Some New Hydantoin Derivatives: Assessment of Activity and Chromatographic Lipophilicity Data

Tatjana Lj. Djaković Sekulić,^{*,a} Adam Smoliński,^b Nemanja P. Trišović,^c
Gordana S. Ušćumlić^c and Biljana Dj. Božić^d

^aDepartment of Chemistry, Biochemistry and Environmental Protection, Faculty of Sciences,
University of Novi Sad, 21000 Novi Sad, Republic of Serbia

^bDepartment of Energy Saving and Air Protection, Central Mining Institute,
40-166 Katowice, Poland

^cFaculty of Technology and Metallurgy and ^dFaculty of Biology, University of Belgrade,
11000 Belgrade, Republic of Serbia

Table S1. Calculated parameters of equation 1: intercept (R_M^0), slope (S) and coefficient of determination (r^2) for mobile phase methanol-water

Compound	R_M^0	S	r^2
1	3.2309	3.9435	0.99519
2	3.1442	3.8590	0.99138
3	2.7644	3.5979	0.96865
4	4.9260	5.8076	0.98996
5	4.1322	4.9649	0.98604
6	4.3032	5.1470	0.97473
7	3.8303	4.8694	0.99305
8	3.8513	4.5542	0.85395
9	4.1122	4.8634	0.90555
10	3.2430	3.9979	0.88405
11	4.3182	4.9498	0.91945
12	4.7313	5.3656	0.90849
13	5.9710	6.5134	0.95304
14	5.2168	5.8810	0.91962
15	4.7547	5.7114	0.95979

*e-mail: tatjana.djakovic-sekulic@dh.uns.ac.rs